

CHRISTIAN COUPPÉ, PT, PhD^{1,4} • RENÉ B. SVENSSON, PhD^{1,3} • KARIN GRÄVARE SILBERNAGEL, PT, ATC, PhD⁵
HENNING LANGBERG, PT, PhD, DSc⁶ • S. PETER MAGNUSSON, PT, DSc^{1,4}

Eccentric or Concentric Exercises for the Treatment of Tendinopathies?

Tendon tissue plays an essential role in transmitting muscle contractile forces to produce movement and is therefore uniquely designed to withstand considerable loads. During locomotion, the Achilles and patellar tendons may see forces up to approximately 8 times body weight.^{23,31,62} Repetitive loading of a tendon often results in overuse injuries, including tendinopathy, which is a clinical condition characterized by pain in the area of the

tendon during activity, localized tenderness on palpation, local swelling of the tendon, and impaired performance.^{45,60} Tendinopathy is a sizable problem in both elite and recreational athletes.^{22,27,93} Specifically, the incidence of tendon injuries has been estimated to be as high as 30% to 50% of all sports injuries and 50% of injuries to elite endurance runners, and 6% of sedentary people will at some point

experience a tendon injury.^{52,58} Moreover, the symptoms and reduction in performance may last for an extended period, potentially years.^{44,57} The exact injury mechanism remains elusive, but an understanding of how tendon tissue adapts to mechanical loading could be the key to understanding the pathogenesis of tendinopathy and thus provide the basis for prevention of overuse injuries.

Tendinopathy is the commonly accepted term for the clinical condition in and around overloaded tendons.⁴⁶ These injuries were previously considered to be the result of a prolonged inflammatory condition (chronic tendinitis).⁹⁰ However, more recently, the extent to which inflammation plays a role in chronic tendinopathy has been debated.^{25,63,80} The slow, insidious onset of tendinopathy makes it difficult to identify the start of the condition in humans, and efforts to develop animal models to study the early events of the condition have been inconsistent.^{32,35} Therefore, the definitive role of inflammation in the early stages of the condition remains difficult to investigate. Notwithstanding these limitations, it has been shown that tendon tissue from individuals with Achilles tendinopathy does not display an elevated expression of inflammatory markers after 1 hour of running.⁷⁷ Rather than inflamed, tendon tissue obtained from individuals with tendinopathy is typically more cellular than healthy tissue and displays both signs of general tissue degeneration, including collagen degeneration and necrosis, as well as signs of regeneration, including neovascularization, irregular fiber structure, and increased ground substance (see Fredberg and Stengaard-Pedersen²⁵).

Tendinopathy is a substantial clinical challenge because it can severely limit sports participation for months and po-

● **SYNOPSIS:** Tendinopathy is a very common disorder in both recreational and elite athletes. Many individuals have recurrent symptoms that lead to chronic conditions and termination of sports activity. Exercise has become a popular and somewhat efficacious treatment regime, and isolated eccentric exercise has been particularly promoted. In this clinical commentary, we review the relevant evidence for different exercise regimes in tendinopathy rehabilitation, with particular focus on the applied loads that are experienced by the tendon and how the exercise regime may affect these applied loads. There is no convincing clinical evidence to demonstrate that isolated eccentric loading exercise improves clinical outcomes more than other loading therapies. However, the great

variation and sometimes insufficient reporting of the details of treatment protocols may hamper the interpretation of what may be the optimal exercise regime with respect to parameters such as load magnitude, speed of movement, and recovery period between exercise sessions. Future studies should control for these loading parameters, evaluate various exercise dosages, and think beyond isolated eccentric exercises to arrive at firm recommendations regarding rehabilitation of individuals with tendinopathies. *J Orthop Sports Phys Ther* 2015;45(11):853-863. Epub 14 Oct 2015. doi:10.2519/jospt.2015.5910

● **KEY WORDS:** *Achilles, forces, load, patellar, recovery, tendon*

¹Institute of Sports Medicine Copenhagen, Bispebjerg Hospital, Copenhagen, Denmark. ²Department of Orthopaedic Surgery, Bispebjerg Hospital, Copenhagen, Denmark. ³Center for Healthy Aging, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark. ⁴Musculoskeletal Rehabilitation Research Unit, Department of Physical Therapy, Bispebjerg Hospital, Copenhagen, Denmark. ⁵Department of Physical Therapy, University of Delaware, Newark, DE. ⁶CopenRehab, Department of Public Health, University of Copenhagen, Copenhagen, Denmark. The authors certify that they have no affiliations with or financial involvement in any organization or entity with a direct financial interest in the subject matter or materials discussed in the article. Address correspondence to Dr S. Peter Magnusson, Institute of Sports Medicine Copenhagen, Bispebjerg Hospital, Bispebjerg Bakke 23, 2400 Copenhagen NV, Denmark. E-mail: p.magnusson@sund.ku.dk ● Copyright ©2015 *Journal of Orthopaedic & Sports Physical Therapy*®




FIGURE 1. Schematic illustration of concentric and eccentric muscle contraction about the ankle joint. Numbers illustrate lengths (not to scale). (A) During concentric heel rise, the tendon and muscle are initially relaxed. (B) As muscle shortens, force is generated, causing the tendon to lengthen until sufficient force has been reached and the heel begins to rise. (C) While muscle shortens further, the heel continues to rise under approximately constant force and, due to the constant load, the length of tendon also remains constant. (D) In the eccentric phase, the heel drops as muscle lengthens, still at approximately constant force, and consequently tendon retains its length. (E) Finally, the muscle lengthens as it is relaxed and tendon shortens because load is removed. Note that across the schematics, tendon length is determined by the amount of load carried independent of muscle length.

tentially years.^{4,44} The list of currently available interventions for this clinical condition is extensive and may include surgery, nonsteroidal anti-inflammatory drugs, corticosteroids, sclerosing injection, shockwave therapy, platelet-rich plasma injection, intratendinous hyperosmolar dextrose (prolotherapy) injection, high-volume injections of 10 mL 0.5% bupivacaine and 40 mL normal saline into the paratenon, Kinesio Tape, and therapeutic ultrasound, just to mention a few.^{81,92} Although these and other treatment options are described in the literature, various loading interventions have predominated in the treatment of tendinopathies.⁶⁴ It seems that loading paradigms yield positive clinical,^{2,61,88} structural,⁵⁰ and biochemical outcomes.⁴⁹

Much of the attention on loading programs as a treatment paradigm for these challenging injuries originated from an article published by Stanish et al⁹⁰ in 1986. In this article, the loading regime was described as a stretch-shortening exercise, that is, an eccentric component rapidly followed by a concentric component. For example, in the case of the patellar tendon, “The patient, from a standing position, flexes the knees and drops to a squatting position abruptly, then recoils to the standing position.”⁹⁰

The progression of the exercise protocol, with abating symptoms, was described as increasing the speed of the movement, and thereafter an external load was added for additional progression.⁹⁰

Approximately a decade later, it was suggested that isolated eccentric contraction alone, without the accompanying concentric component of a stretch-shortening cycle, provided good clinical results for patients with tendinopathy.⁴ This isolated eccentric loading paradigm has since gained considerable popularity and is now widely regarded as the treatment of choice, although there is a lack of convincing evidence that it is the most effective exercise regime.⁶⁴ More recently, new loading-based exercise regimes, such as isolated concentric training,⁶¹ heavy slow resistance training,^{9,49} and concentric/eccentric progressing to eccentric training,^{88,89} have emerged. This article focuses on the underlying rationale of these various loading paradigms.

Response of Healthy Tendon to Load

It is well established that exercise in general can affect both skeletal muscle³⁸ and tendon.⁴⁸ In tendon, there is an acute increase in blood flow and collagen synthesis,^{54,55} and long-term effects lead to tissue hypertrophy and altered material

properties.^{12,85,98} The magnitude and type of adaptation likely depend on the exercise regime, including the magnitude of the load, range of motion performed, contraction mode (eccentric lengthening/concentric shortening), movement speed, number of repetitions, and rest periods between the exercise sessions. By varying these components, a wide range of exercise programs can be constructed, from endurance (low load, high speed, many repetitions) to strength (high load, low speed, few repetitions) programs, with myriad combinations in between. The response of tendon to the various exercise parameters will be discussed in the following sections, though knowledge on several matters remains limited.

It is well known that the tendon cells (fibroblasts) respond to mechanical stimuli in the form of strain,^{42,66,96} and that depriving them of strain (relative tissue deformation) leads to degeneration and apoptosis (cell death).^{7,17,99} However, the dose response to strain magnitude is still not well established. Cell-culture experiments suggest that there is an increased response (increased collagen expression, reduced matrix metalloproteinase expression, and increased matrix stiffness) with increased strain,^{56,101} but there may be an optimal strain, beyond which the

stretching becomes detrimental.^{41,68} The absolute values of strain used in cell-culture stimulation vary greatly, and it is uncertain how much of a given strain the cells will experience in vivo, because the surrounding matrix may provide shielding. In vivo, in healthy human Achilles tendons, it has been reported that with equal exercise volume, working at 90% of maximum voluntary contraction (MVC), which causes approximately 5% tendon strain, yields increased stiffness and cross-sectional area compared to working at 55% MVC, which causes approximately 3% tendon strain.⁵

Speed and/or duration of loading during exercise also appear to be important for tendon adaptation.^{6,41,51,56} At the cellular level, most,^{30,41,97} but not all,²¹ studies have found that the adaptive response of fibroblasts to dynamic load is superior to that of static load (zero speed). Though the response to different dynamic load regimes is complex due to the interaction of the parameters, overall, the evidence suggests that increased time under load, increased number of load cycles, and increased loading rate result in a positive adaptive response (increased matrix strength and stiffness and decreased matrix metalloproteinase expression) in cultured fibroblasts.^{41,56,100} The response appears to be a bit different in vivo, where it has been reported that the stiffness and size (cross-sectional area) of human Achilles tendon were more responsive to a low number of loads of long duration (6-second cycle) than to a high number of faster loads (2-second cycle) when the total exercise volume was kept constant.^{5,6} This finding is corroborated somewhat by another study showing that isometric contractions of long duration (20 seconds) yielded greater patellar tendon adaptation (increased stiffness) than rapid (1-second) contractions at equal exercise volume.⁵¹ In the latter study, muscle strength and volume adaptation were unaffected by contraction duration.

It is unknown whether tendon cells experience some form of fatigue as a result of repeated load cycles. Tendon cells

have a low metabolic rate and probably do not require rest for the purpose of restoring energy deposits. However, the anabolic response to loading is sustained in tendon for up to several days following an exercise bout,^{55,65} which could indicate the need for a postexercise recovery period.⁶³ But, conversely, studies on cell culture have performed continuous stimulation of fibroblasts for up to 24 hours per day without detrimental effects,^{41,59} and most exercise protocols for tendinopathy management are performed every day without recovery periods.^{4,67,89} Overall, there is a lack of studies specifically addressing the effects of recovery and how it affects tendon adaptation.

The majority of the tendon is composed of extracellular matrix (ECM), which is a passive structure and, unlike the cells, does not actively respond to load, though it may still be differentially affected by exercise parameters. It has been suggested that accumulation of microdamage may be involved in the etiology of tendinopathy⁸⁷ and, because the turnover in tendons is slow,³⁴ ECM damage could accumulate. Microdamage is difficult to measure, and its clinical relevance is therefore unclear. However, if it does play a role, it would be an argument in favor of recovery periods. Mechanical studies have shown that both overloading and mechanical fatigue can cause damage to tendon ECM,^{43,72} which could play a role in tendinopathy. The tendon ECM is also a viscoelastic material, which means that slower loading regimes can yield greater strains than faster loading regimes, as the tendon has more time to creep.⁷⁴ Creep also appears to be associated with greater relative fibril slippage,²⁴ which may generate local shear strains sensed by the cells. Slow loading may therefore produce particularly strong cell stimuli that can be beneficial to the tendon if the strain is sufficient, but could be detrimental if strain is excessive. This viscoelastic behavior depends on the amount of time the tendon is under load and is therefore unaffected by the mode of muscle contraction (eccentric or concentric).




FIGURE 2. Achilles tendon force versus Achilles tendon length. Eccentric data are shown as open circles, whereas concentric data are filled circles. Reprinted with permission from Chaudhry S, Morrissey D, Woledge RC, Bader DL, Screen HR. Eccentric and concentric loading of the triceps surae: an in vivo study of dynamic muscle and tendon biomechanical parameters. *J Appl Biomech.* 2015;31:69-78. ©Human Kinetics, Inc.

In summary, tendon is responsive to loading and will respond more strongly to greater loads, although there is likely an optimum beyond which load becomes detrimental. Slower loading regimes may be superior to rapid loading, while the importance of recovery between loading sessions is unclear.

Tendon Under Eccentric and Concentric Muscle Contractions

Although isolated eccentric loading regimes for tendinopathy have been widely accepted as the treatment of choice,⁹² the potential mechanisms behind this intervention remain unclear. In the following section, we discuss some of the proposed mechanisms and their potential applications in light of existing evidence.

Strictly speaking, the descriptions “concentric” and “eccentric” only apply to muscle, which actively contracts. Tendon is a mechanically passive structure that lengthens when load increases and shortens when load is reduced (**FIGURE 1**). It is therefore questionable whether the mode of muscle contraction for a given load and range of motion would have a differential effect on tendon tissue. The fact that

muscles can produce greater maximal force eccentrically than concentrically would suggest a potential for greater mechanical stimulation from eccentric than from concentric exercise.^{18,37} Accordingly, it has been suggested that the tendon may stretch more during eccentric loading than concentric loading.^{1,90} However, while there is a potential for greater tendon load and, consequently, stretch with eccentric exercise, this potential is rarely utilized because rehabilitation exercises seldom approach concentric 1-repetition maximum. It has been shown that Achilles tendon load and stretch are identical during the concentric and eccentric components of a heel rise/drop against body weight (FIGURE 2), a typical load used in rehabilitation.^{11,79} The 2 most commonly used eccentric exercises for tendinopathy are squats for the patellar tendon and standing heel lowering for the Achilles tendon, and both movements are typically performed to a 15-repetition maximum.^{4,103}

TABLES 1 and 2 list the number of weekly repetitions and estimated loads used in several eccentric exercise regimes published in the literature. It should be noted that several studies have used an individualized load progression, but they seldom report the actual amount achieved, only the starting load. It is also worth noting that most studies do not include a control group, which is likely due to ethical concerns; therefore, it is uncertain whether the provided treatment was in fact superior to a “wait and see” approach. However, most of these studies included patients with chronic symptoms, suggesting that time alone would not have improved their condition.

A few theoretical mechanisms have been suggested to explain how eccentric loading may differentially influence tendon. For example, during heel-lowering/raising tasks, the ground reaction force may fluctuate at a higher frequency (approximately 10 Hz) during the eccentric phase.^{36,79,84} While tendon cells could potentially register this modulation, the magnitude of the modulation is quite

TABLE 1		
EXERCISE PROGRAMS USED IN STUDIES FOR ACHILLES TENDINOPATHY		
Study/Group	Load*	Pain Improvement†
Alfredson et al ⁴		
Isolated ECC	1260 (100%) [§]	VAS, 94%
Croisier et al ¹³		
Isolated ECC	135 (130%)	VAS, 73% (10 wk)
Alfredson and Lorentzon ³		
Isolated ECC	1260 (100%) [§]	VAS, 75%
Öhberg and Alfredson ⁷⁰		
Isolated ECC	1260 (100%) [§]	Pain during activity: reduced, 88%
Roos et al ⁸³		
Isolated ECC	1260 (100%) [§]	Foot and Ankle Outcome Score, 36%
Shalabi et al ⁸⁶		
Isolated ECC	1260 (100%) [§]	6-point pain scale, 40%
de Vos et al ¹⁶		
Isolated ECC	1260 (100%) [§]	VISA-A, 78%
Langberg et al ⁵³		
Isolated ECC	1260 (120%) [§]	VAS, 71%
Petersen et al ⁷⁶		
Isolated ECC	1890 (100%) [§]	VAS, 60%
Yelland et al ¹⁰²		
Isolated ECC	1260 (100%) [§]	VISA-A, 38%
Silbernagel et al ⁸⁸		
Isolated ECC plus mixed ECC + CON plus other exercises	Mixed, 840 (50%) [‡] Mixed, 945 (100%) [§] ECC, 630 (100%) [§]	VISA-A, 60%
Rompe et al ⁸²		
Isolated ECC	1260 (100%) [§]	VISA-A, 49% (16 wk) [¶]
Wait and see	0 (0%)	VISA-A, 14% (16 wk)

Table continues on page 857.

small compared to the total load (approximately 10%). It has also been proposed that motor unit activation differs between concentric and eccentric exercises,¹⁸ which may produce a difference in load distribution and shear within the tendon, although such a mechanism is unlikely to be a factor for the patellar tendon, in which load is distributed via the patella. Finally, secreted signaling molecules from the muscle (myokines), such as interleukin 6 (IL-6),⁷⁵ could potentially affect the nearby tendon, and myokine expression may differ between concen-

tric and eccentric exercise, although the effects are currently unclear.^{73,94}

In contrast to these hypothetical considerations, there is evidence to suggest that eccentric and concentric muscle contractions do not yield a differential response of the tendon. In animal models, it has been shown that concentric or eccentric contraction to the same force level does not influence the expression of collagen at the cellular level.²⁹ In fact, even if the eccentric muscular contraction force were greater than concentric contraction force, both contraction modes would yield

TABLE 1

EXERCISE PROGRAMS USED IN STUDIES
FOR ACHILLES TENDINOPATHY (CONTINUED)

Study/Group	Load*	Pain Improvement†
Nørregaard et al ⁶⁹		
Isolated ECC	1260 (100%) ^{‡§}	Modified KOOS-score: improved, 25%
Stretching	Stretching	Modified KOOS-score: improved, 25%
Beyer et al ⁹		
Isolated ECC	1260 (100%) ^{‡§}	VISA-A, 24% VAS, 59%
Heavy slow resistance (ECC + CON)	144 (130%) [‡]	VISA-A, 41% VAS, 69%
Mafi et al ⁶¹		
Isolated ECC	1260 (100%) ^{‡§}	Satisfied, 82% [¶] VAS of satisfied, 83%
Isolated CON plus jump plus rope skipping	1260 (100%) [‡]	Satisfied, 32% VAS of satisfied, 86%
Niesen-Vertommen et al ⁶⁷		
Isolated ECC	300 (55%) [§]	VAS, 78% [¶]
Mixed ECC + CON	300 (25%) ^{¶¶}	VAS, 46%
Silbernagel et al ⁸⁹		
Isolated ECC plus mixed ECC + CON plus other exercises	Mixed, 840 (50%) [‡] Mixed, 945 (100%) ^{‡§} ECC, 630 (100%) ^{‡§}	Pain during activity: reduced, 57%
Mixed ECC + CON	1260 (50%) or 315 (100%) [§]	Pain during activity: reduced, 80%

Abbreviations: CON, concentric; ECC, eccentric; KOOS, Knee injury Osteoarthritis Outcome Score; VAS, visual analog scale; VISA-A, Victorian Institute of Sport Assessment-Achilles.

*Load reported as weekly repetitions and estimated load as percentage of body weight (50% body weight is 2 legged and 100% is 1 legged). In general, peak load is listed, but for studies with load progression that did not report the achieved progress, only the load before progression is listed. No attempt was made to evaluate load in stretching exercises.

†Unless otherwise stated, all studies had 12 weeks of intervention.

‡Pain accepted.

§Study included unreported progression beyond what is listed.

¶Estimated percent body weight from maximal voluntary contraction or other external load.

¶¶Improvement in ECC group significantly greater than in comparator.

similar expressions of collagen.³³ These animal findings imply that given a sufficiently high force (and resulting strain on the fibroblast), the contraction mode is inconsequential for the tendon cellular response. A recent study examined the effect of contraction mode on tendon (and muscle) hypertrophy in healthy human subjects.²⁰ The 12-week resistance training consisted of isolated concentric knee extensions on one side and eccentric knee extensions on the contralateral side. The sets, repetitions, and time of loading were similar between sides, but the loading for the eccentric side was 120%

that of the concentric side. The results showed that resistance training with either concentric or eccentric contraction produced a similar magnitude of tendon hypertrophy.²⁰ These findings reinforce the notion that the cellular and tissue response in healthy tendon is independent of contraction mode.

In summary, there are a number of mechanisms that could theoretically differentiate the effect of eccentric from concentric exercise on tendon, but there is no evidence that these mechanisms actually play a role or are beneficial. In contrast, there is evidence from animal

and human studies that suggests a lack of differential effect of eccentric versus concentric exercises.

The Effects of Muscle Contraction Regimes on Tendinopathy

Both elite and recreational athletes are frequently afflicted by Achilles and/or patellar tendinopathy.^{52,57} The clinical literature has mainly focused on use of eccentric loading exercise in the treatment of these tendinopathies, and therefore the following sections on Achilles and patellar tendinopathy will first cover the clinical studies that have investigated the effects of eccentric loading, followed by studies that have investigated the effects of other types of loading exercises. TABLES 1 and 2 cover the clinical studies on Achilles and patellar tendinopathy, with load reported as weekly repetitions and estimated load as a percentage of body weight.

The Effect of Eccentric Muscle Contraction on Achilles Tendon It has been shown that eccentric-loading regimens for Achilles tendinopathy can provide clinical improvements, including reduction in pain and improved function.^{15,47,88,92} Several studies^{3,13} have employed the isolated eccentric-loading paradigm initially introduced by Alfredson et al.⁴ When this exercise model is performed as unilateral heel drop, the force placed on the tendon is a function of body weight, and the force can be modulated with additional weight placed in a backpack worn by the patient. In addition to improvements in pain and function, it appears that structural features observed with ultrasound and magnetic resonance imaging are altered following isolated eccentric loading in some,^{28,70,71,86} but not all, studies.^{14,69,76,82} It has also been shown that in addition to decreasing pain, isolated eccentric loading can result in increased synthesis of type I collagen.⁵³ Thus, isolated eccentric loading appears to influence biochemical and biomechanical parameters and improve clinical outcomes. The beneficial effects of isolated eccentric loading with body weight appear to be reduced if the pain is located toward the tendon inser-

tion¹⁹; however, this might be remedied by avoiding ankle dorsiflexion below horizontal to avoid compression of the distal end of the tendon against the posterior aspect of the calcaneus.⁴⁰

Several studies have compared isolated eccentric loading with body weight to other types of nonloading therapy (eg, prolotherapy, cryotherapy, splints).^{16,76,83,102} Most of these studies report significant clinical improvements with eccentric exercise,^{16,76,83,102} although the effect relative to the alternative treatment varies (eg, eccentric loading regimes have greater effect than cryotherapy, but similar effect compared to shockwave therapy and heel brace).⁹²

Collectively, these studies demonstrate positive clinical benefits from eccentric loading, but because none of these studies has a comparison group using an alternative muscle contraction mode (concentric or isometric), they are unable to show whether the actual muscle contraction mode plays a role in the outcome.

The Effect of Eccentric Muscle Contraction Compared to Other Contraction Regimes for Achilles Tendinopathy A limited number of studies have investigated the effect of isolated eccentric exercise relative to other loading regimes for Achilles tendinopathy.^{9,61,67,69,89} It has been shown that fast heel-drop exercises with a high eccentric load were more effective in reducing overall pain than isotonic concentric/eccentric exercises at lower loads.⁶⁷ Others have compared isolated eccentric and concentric loading and demonstrated similar pain reduction with both regimes, with the load on the tendon being greater in the eccentric (body weight plus backpack) than in the concentric (less or equal body weight) group (TABLE 1).⁶¹ A more extensive rehabilitation program that included concentric/eccentric loading with body weight yielded better self-reported outcomes (questionnaire on function and pain) than a concentric program of a lesser total loading volume not at 6 weeks but at 1-year follow-up.⁸⁹ Isolated eccentric loading with body weight has also been compared to static

TABLE 2		EXERCISE PROGRAMS USED IN STUDIES FOR PATELLAR TENDINOPATHY	
Study/Group	Load*	Pain Improvement†	
Bahr et al ⁸			
Isolated ECC	630 (100%) [§]	VISA-P, 73%	
Croisier et al ¹³			
Isolated ECC	135 (130%) [¶]	VAS, 71%	
Purdam et al ⁷⁸			
Isolated ECC	630 (100%) [§]	VAS, 62%	
Stasinopoulos and Stasinopoulos ⁹¹			
Isolated ECC	135 (100%) [§]	5-point scale, 80% (4 wk)	
Visnes et al ⁹⁵			
Isolated ECC + competitive volleyball	630 (100%) [§]	No improvement	
Competitive volleyball (control)	0 (0%)	No improvement	
Young et al ¹⁰³			
Isolated ECC on 25° decline board	630 (100%) [§]	VISA-P, 25% VAS, 51%	
Isolated ECC on horizontal step	630 (100%)	VISA-P, 18% VAS, 53%	
Jonsson and Alfredson ³⁹			
Isolated ECC	630 (100%) [§]	VISA-P, 102% [¶] VAS, 69% [¶]	
Isolated CON (n = 4)	630 (100%) [§]	VISA-P, -9% VAS, 8%	

Table continues on page 859.

stretching, which places a low-magnitude load on the tendon for an extended period, and the results showed that the interventions yielded a similar clinical outcome.⁶⁹ It was recently demonstrated that heavy slow resistance training 3 times per week was equally effective in reducing symptoms compared to the traditional eccentric regimen performed 7 days per week in patients with Achilles tendinopathy.⁹ These data indicate that different muscle contraction loading regimes can accomplish the same clinical improvement in patients with Achilles tendinopathy. Thus, it seems that loading confers some clinical improvements to Achilles tendinopathy, but, based on the available literature, it is not possible to delineate the role of contraction mode (eccentric or concentric) from that of load magnitude, number of repetitions and sets, contraction speed, and recovery time between sessions.

The Effects of Eccentric Muscle Contraction on Patellar Tendon Similar to Achilles tendinopathy, it has been suggested that eccentric-loading regimes for patellar tendinopathy may provide clinical improvements.^{8,13,39,78} Patients typically eccentrically load the affected patellar tendon while performing a partial squat on the affected limb, and then return to the starting position by concentrically loading the nonaffected tendon. One study compared isolated eccentric loading with body weight to nonloading therapy with ultrasound or transverse friction massage.⁹¹ The authors reported that as few as 3 weekly sessions (each session consisting of 3 sets of 15 repetitions) of isolated eccentric squats for 4 weeks yielded significant clinical improvements, which were far greater than with nonloading therapy.⁹¹ It has also been shown that eccentric loading can result

TABLE 2

EXERCISE PROGRAMS USED IN STUDIES
FOR PATELLAR TENDINOPATHY (CONTINUED)

Study/Group	Load*	Pain Improvement†
Kongsgaard et al ⁴⁹		
Isolated ECC	630 (100%)‡	VISA-P, 42%
Heavy slow resistance (ECC + CON)	288 (130%)‡	VISA-P, 39%
Kongsgaard et al ⁵⁰		
No ECC
Heavy slow resistance (ECC + CON)	288 (130%)‡	VISA-P, 27%
Frohm et al ²⁶		
Isolated ECC	315 (100%)§	VISA-P, 108%
Isolated ECC overload	32 (170%)¶	VISA-P, 76%
Cannell et al ¹⁰		
ECC drop squats	300 (60%)§	VAS, 55%
Mixed ECC + CON, focused on CON	150 (CON, 125%)§¶ (ECC, 63%)¶	VAS, 31%

Abbreviations: CON, concentric; ECC, eccentric; VAS, visual analog scale; VISA-P, Victorian Institute of Sport Assessment-patella.

*Load reported as weekly repetitions and estimated load as percentage of body weight (50% body weight is 2 legged and 100% is 1 legged). In general, peak load is listed, but for studies with load progression that did not report the achieved progress, only the load before progression is listed. No attempt was made to evaluate load in stretching exercises.

†Unless otherwise stated, all studies had 12 weeks of intervention.

‡Pain accepted.

§Study included unreported progression beyond what is listed.

¶Estimated percent body weight from maximal voluntary contraction or other external load.

¶Improvement in ECC group significantly greater than in comparator.

pants remained in the concentric group, hampering any meaningful comparison and firm conclusion.

Other studies have compared loading regimes without matching the load (TABLE 2). Isolated eccentric squats on a decline board have been compared to mixed concentric/eccentric squats on a flat surface in elite volleyball players: both groups performed exercises on only the affected extremity for 12 weeks before starting the competitive season.¹⁰³ Both protocols resulted in reduced pain and increased function, without a difference between the groups. Twelve weeks of isolated heavy eccentric training using few repetitions of high load (100% MVC, approximately 170% body weight) has been compared with isolated eccentric squats on a decline board using a larger number of repetitions at lower load (100% body weight, approximately 60% MVC).²⁶ Both groups markedly improved in pain and function but without a difference between the groups. Another study compared eccentric drop squats landing on 2 feet (60 repetitions per session) with knee flexion/extension exercise (30 repetitions per session), with the aim of creating a high concentric load component (75%-80% MVC).¹⁰ Both groups trained with progressive loads 5 times weekly and had reduced tendon pain after 12 weeks, but outcomes were no different between groups. Only 1 study has investigated biochemical outcomes: the efficacy of isolated eccentric squats compared to mixed concentric/eccentric heavy slow resistance training.⁴⁹ Both exercise regimes resulted in reduced pain and improved function, but biochemical changes (increased collagen content and reduced glycation) were only evident with heavy slow resistance training. In addition, heavy slow resistance training was associated with structural changes.⁵⁰ Therefore, collectively, there is no firm evidence to support the notion that eccentric loading is more efficient than concentric or other loading regimes for patellar tendinopathy. Most studies have not matched for other parameters, such

in clinical improvement when patients were instructed to refrain from sports activity during the intervention period.^{39,91} However, in elite volleyball players with patellar tendinopathy, adding 8 weeks of isolated eccentric loading to the already existing activity during the season did not confer any pain relief.⁹⁵ In contrast, in Achilles tendinopathy, it has been shown that moderate physical activity during the treatment period was not detrimental to the benefits of eccentric exercise.⁸⁸ The apparent difference between activity at an elite level and more moderate activity may be related to insufficient recovery time or possibly the total load volume of sports activity and the added eccentric-loading exercises,⁶³ although this requires further investigation. Ultrasonography has been used to investigate structural changes (tendon thickness and Doppler activity) in patellar tendinopa-

thy following eccentric exercise, but has not demonstrated significant effects.⁴⁹ Collectively, the above studies show that eccentric loading may provide clinical benefits in patellar tendinopathy treatment, but, again, whether the actual direction of muscle contraction plays a role in the outcome or clinical benefits are instead related to the absolute magnitude of loading is unclear, as none of these studies had a similar comparison group. **The Effect of Eccentric Muscle Contraction Compared to Other Contraction Regimes for Patellar Tendinopathy** For patellar tendinopathy, only 1 study has compared isolated eccentric loading with an identical concentric-loading regime at equal load magnitude, volume, and speed.³⁹ The authors³⁹ concluded that eccentric loading was more effective in reducing pain than concentric loading; however, due to a high number of dropouts, only 4 of 7 partici-

as load, speed, frequency, and rest periods,^{10,26,39} and it is therefore still unclear what specific load magnitude, frequency, or total load volume per session should be used to provide meaningful clinical and structural improvements in patients with patellar tendinopathy.

Clinical Implications and Future Directions

Tendinopathies are a common ailment among athletes, and there is currently an incomplete picture of the etiology. Although there are several suggested treatment options, loading regimes appear to have good clinical results. Early works have shown an exercise program with isolated eccentric contraction to have good clinical outcomes, which promoted the paradigm of eccentric loading for tendinopathy. However, as outlined in this review, there is little evidence for isolating the eccentric component of a loading-based regime. The basic mechanisms that are likely to influence tendon adaptations appear to be related mainly to tendon load/strain magnitude and duration, and there is no theoretical basis for greater tendon loads in eccentric exercises at a given force (body weight or external load).

There is a paucity of clinical trials directly comparing different exercise regimes and different exercise dosages, but the available evidence provides little support for the superiority of isolated eccentrics. It is worth noting that studies rarely use comparable load magnitude when comparing eccentric to other load regimes. To delineate the effects of mode and load magnitude, future work should compare isolated eccentric and concentric action under equal load at various exercise dosages in individuals with tendinopathy. The focus on eccentric exercise has overshadowed other aspects of tendinopathy rehabilitation, and acknowledgement of the limited evidence may prompt a broader approach, including the use of heavy load and low speed, which has some support from both basic science and clinical trials. ●

REFERENCES

- Alfredson H. The chronic painful Achilles and patellar tendon: research on basic biology and treatment. *Scand J Med Sci Sports*. 2005;15:252-259. <http://dx.doi.org/10.1111/j.1600-0838.2005.00466.x>
- Alfredson H, Lorentzon R. Chronic Achilles tendinosis: recommendations for treatment and prevention. *Sports Med*. 2000;29:135-146.
- Alfredson H, Lorentzon R. Intratendinous glutamate levels and eccentric training in chronic Achilles tendinosis: a prospective study using microdialysis technique. *Knee Surg Sports Traumatol Arthrosc*. 2003;11:196-199. <http://dx.doi.org/10.1007/s00167-003-0360-0>
- Alfredson H, Pietilä T, Jonsson P, Lorentzon R. Heavy-load eccentric calf muscle training for the treatment of chronic Achilles tendinosis. *Am J Sports Med*. 1998;26:360-366.
- Arampatzis A, Karamanidis K, Albracht K. Adaptational responses of the human Achilles tendon by modulation of the applied cyclic strain magnitude. *J Exp Biol*. 2007;210:2743-2753. <http://dx.doi.org/10.1242/jeb.003814>
- Arampatzis A, Peper A, Bierbaum S, Albracht K. Plasticity of human Achilles tendon mechanical and morphological properties in response to cyclic strain. *J Biomech*. 2010;43:3073-3079. <http://dx.doi.org/10.1016/j.jbiomech.2010.08.014>
- Arnoczky SP, Lavagnino M, Egerbacher M. The mechanobiological aetiopathogenesis of tendinopathy: is it the over-stimulation or the under-stimulation of tendon cells? *Int J Exp Pathol*. 2007;88:217-226. <http://dx.doi.org/10.1111/j.1365-2613.2007.00548.x>
- Bahr R, Fossan B, Løken S, Engebretsen L. Surgical treatment compared with eccentric training for patellar tendinopathy (jumper's knee). A randomized, controlled trial. *J Bone Joint Surg Am*. 2006;88:1689-1698. <http://dx.doi.org/10.2106/JBJS.E.01181>
- Beyer R, Kongsgaard M, Hougs Kjaer B, Øhlenschläger T, Kjaer M, Magnusson SP. Heavy slow resistance versus eccentric training as treatment for Achilles tendinopathy: a randomized controlled trial. *Am J Sports Med*. 2015;43:1704-1711. <http://dx.doi.org/10.1177/0363546515584760>
- Cannell LJ, Taunton JE, Clement DB, Smith C, Khan KM. A randomised clinical trial of the efficacy of drop squats or leg extension/leg curl exercises to treat clinically diagnosed jumper's knee in athletes: pilot study. *Br J Sports Med*. 2001;35:60-64. <http://dx.doi.org/10.1136/bjism.35.1.60>
- Chaudhry S, Morrissey D, Woledge RC, Bader DL, Screen HR. Eccentric and concentric loading of the triceps surae: an in vivo study of dynamic muscle and tendon biomechanical parameters. *J Appl Biomech*. 2015;31:69-78.
- Couppé C, Kongsgaard M, Aagaard P, et al. Habitual loading results in tendon hypertrophy and increased stiffness of the human patellar tendon. *J Appl Physiol* (1985). 2008;105:805-810. <http://dx.doi.org/10.1152/japplphysiol.90361.2008>
- Croisier JL, Forthomme B, Foidart-Dessalle M, Godon B, Crielaard JM. Treatment of recurrent tendinitis by isokinetic eccentric exercises. *Iso-kinet Exerc Sci*. 2001;9:133-141.
- de Jonge S, de Vos RJ, Van Schie HT, Verhaar JA, Weir A, Tol JL. One-year follow-up of a randomised controlled trial on added splinting to eccentric exercises in chronic midportion Achilles tendinopathy. *Br J Sports Med*. 2010;44:673-677. <http://dx.doi.org/10.1136/bjism.2008.052142>
- de Vos RJ, Weir A, van Schie HT, et al. Platelet-rich plasma injection for chronic Achilles tendinopathy: a randomized controlled trial. *JAMA*. 2010;303:144-149. <http://dx.doi.org/10.1001/jama.2009.1986>
- de Vos RJ, Weir A, Visser RJ, de Winter T, Tol JL. The additional value of a night splint to eccentric exercises in chronic midportion Achilles tendinopathy: a randomised controlled trial. *Br J Sports Med*. 2007;41:e5. <http://dx.doi.org/10.1136/bjism.2006.032532>
- Dideriksen K. Muscle and tendon connective tissue adaptation to unloading, exercise and NSAID. *Connect Tissue Res*. 2014;55:61-70. <http://dx.doi.org/10.3109/03008207.2013.862527>
- Enoka RM. Eccentric contractions require unique activation strategies by the nervous system. *J Appl Physiol* (1985). 1996;81:2339-2346.
- Fahlström M, Jonsson P, Lorentzon R, Alfredson H. Chronic Achilles tendon pain treated with eccentric calf-muscle training. *Knee Surg Sports Traumatol Arthrosc*. 2003;11:327-333. <http://dx.doi.org/10.1007/s00167-003-0418-z>
- Farup J, Rahbek SK, Vendelbo MH, et al. Whey protein hydrolysate augments tendon and muscle hypertrophy independent of resistance exercise contraction mode. *Scand J Med Sci Sports*. 2014;24:788-798. <http://dx.doi.org/10.1111/sms.12083>
- Feng Z, Tateishi Y, Nomura Y, Kitajima T, Nakamura T. Construction of fibroblast-collagen gels with orientated fibrils induced by static or dynamic stress: toward the fabrication of small tendon grafts. *J Artif Organs*. 2006;9:220-225. <http://dx.doi.org/10.1007/s10047-006-0354-z>
- Ferretti A. Epidemiology of jumper's knee. *Sports Med*. 1986;3:289-295.
- Finni T, Komi PV, Lepola V. In vivo human triceps surae and quadriceps femoris muscle function in a squat jump and counter movement jump. *Eur J Appl Physiol*. 2000;83:416-426. <http://dx.doi.org/10.1007/s004210000289>
- Folkhard W, Geercken W, Knörzer E, et al. Structural dynamic of native tendon collagen. *J Mol Biol*. 1987;193:405-407. [http://dx.doi.org/10.1016/0022-2836\(87\)90228-2](http://dx.doi.org/10.1016/0022-2836(87)90228-2)
- Fredberg U, Stengaard-Pedersen K. Chronic tendinopathy tissue pathology, pain mechanisms, and etiology with a spe-

- cial focus on inflammation. *Scand J Med Sci Sports*. 2008;18:3-15. <http://dx.doi.org/10.1111/j.1600-0838.2007.00746.x>
26. Frohm A, Saartok T, Halvorsen K, Renström P. Eccentric treatment for patellar tendinopathy: a prospective randomised short-term pilot study of two rehabilitation protocols. *Br J Sports Med*. 2007;41:e7. <http://dx.doi.org/10.1136/bjism.2006.032599>
27. Frost P, Bonde JP, Mikkelsen S, et al. Risk of shoulder tendinitis in relation to shoulder loads in monotonous repetitive work. *Am J Ind Med*. 2002;41:11-18. <http://dx.doi.org/10.1002/ajim.10019>
28. Gårdin A, Bruno J, Movin T, Kristoffersen-Wiberg M, Shalabi A. Magnetic resonance signal, rather than tendon volume, correlates to pain and functional impairment in chronic Achilles tendinopathy. *Acta Radiol*. 2006;47:718-724. <http://dx.doi.org/10.1080/02841850600774035>
29. Garma T, Kobayashi C, Haddad F, Adams GR, Bodell PW, Baldwin KM. Similar acute molecular responses to equivalent volumes of isometric, lengthening, or shortening mode resistance exercise. *J Appl Physiol* (1985). 2007;102:135-143. <http://dx.doi.org/10.1152/japplphysiol.00776.2006>
30. Gauvin R, Parenteau-Bareil R, Larouche D, et al. Dynamic mechanical stimulations induce anisotropy and improve the tensile properties of engineered tissues produced without exogenous scaffolding. *Acta Biomater*. 2011;7:3294-3301. <http://dx.doi.org/10.1016/j.actbio.2011.05.034>
31. Giddings VL, Beaupré GS, Whalen RT, Carter DR. Calcaneal loading during walking and running. *Med Sci Sports Exerc*. 2000;32:627-634.
32. Glazebrook MA, Wright JR, Jr., Langman M, Stanish WD, Lee JM. Histological analysis of Achilles tendons in an overuse rat model. *J Orthop Res*. 2008;26:840-846. <http://dx.doi.org/10.1002/jor.20546>
33. Heinemeier KM, Olesen JL, Haddad F, et al. Expression of collagen and related growth factors in rat tendon and skeletal muscle in response to specific contraction types. *J Physiol*. 2007;582:1303-1316. <http://dx.doi.org/10.1113/jphysiol.2007.127639>
34. Heinemeier KM, Schjerling P, Heinemeier J, Magnusson SP, Kjaer M. Lack of tissue renewal in human adult Achilles tendon is revealed by nuclear bomb ¹³C. *FASEB J*. 2013;27:2074-2079. <http://dx.doi.org/10.1096/fj.12-225599>
35. Heinemeier KM, Skovgaard D, Bayer ML, et al. Uphill running improves rat Achilles tendon tissue mechanical properties and alters gene expression without inducing pathological changes. *J Appl Physiol* (1985). 2012;113:827-836. <http://dx.doi.org/10.1152/japplphysiol.00401.2012>
36. Henriksen M, Aaboe J, Bliddal H, Langberg H. Biomechanical characteristics of the eccentric Achilles tendon exercise. *J Biomech*. 2009;42:2702-2707. <http://dx.doi.org/10.1016/j.jbiomech.2009.08.009>
37. Hill AV. The heat of shortening and the dynamic constants of muscle. *Proc R Soc Lond B Biol Sci*. 1938;126:136-195. <http://dx.doi.org/10.1098/rspb.1938.0050>
38. Holloszy JO, Coyle EF. Adaptations of skeletal muscle to endurance exercise and their metabolic consequences. *J Appl Physiol Respir Environ Exerc Physiol*. 1984;56:831-838.
39. Jonsson P, Alfredson H. Superior results with eccentric compared to concentric quadriceps training in patients with jumper's knee: a prospective randomised study. *Br J Sports Med*. 2005;39:847-850. <http://dx.doi.org/10.1136/bjism.2005.018630>
40. Jonsson P, Alfredson H, Sunding K, Fahlström M, Cook J. New regimen for eccentric calf-muscle training in patients with chronic insertional Achilles tendinopathy: results of a pilot study. *Br J Sports Med*. 2008;42:746-749. <http://dx.doi.org/10.1136/bjism.2007.039545>
41. Joshi SD, Webb K. Variation of cyclic strain parameters regulates development of elastic modulus in fibroblast/substrate constructs. *J Orthop Res*. 2008;26:1105-1113. <http://dx.doi.org/10.1002/jor.20626>
42. Kalson NS, Holmes DF, Herchenhan A, Lu Y, Starborg T, Kadler KE. Slow stretching that mimics embryonic growth rate stimulates structural and mechanical development of tendon-like tissue in vitro. *Dev Dyn*. 2011;240:2520-2528. <http://dx.doi.org/10.1002/dvdy.22760>
43. Ker RF, Wang XT, Pike AV. Fatigue quality of mammalian tendons. *J Exp Biol*. 2000;203:1317-1327.
44. Kettunen JA, Kvist M, Alanen E, Kujala UM. Long-term prognosis for jumper's knee in male athletes. A prospective follow-up study. *Am J Sports Med*. 2002;30:689-692.
45. Khan K, Cook J. The painful nonruptured tendon: clinical aspects. *Clin Sports Med*. 2003;22:711-725.
46. Khan KM, Cook JL, Kannus P, Maffulli N, Bonar SF. Time to abandon the "tendinitis" myth. *BMJ*. 2002;324:626-627. <http://dx.doi.org/10.1136/bmj.324.7338.626>
47. Kingma JJ, de Knikker R, Wittink HM, Takken T. Eccentric overload training in patients with chronic Achilles tendinopathy: a systematic review. *Br J Sports Med*. 2007;41:e3. <http://dx.doi.org/10.1136/bjism.2006.030916>
48. Kjaer M. Role of extracellular matrix in adaptation of tendon and skeletal muscle to mechanical loading. *Physiol Rev*. 2004;84:649-698. <http://dx.doi.org/10.1152/physrev.00031.2003>
49. Kongsgaard M, Kovanen V, Aagaard P, et al. Corticosteroid injections, eccentric decline squat training and heavy slow resistance training in patellar tendinopathy. *Scand J Med Sci Sports*. 2009;19:790-802. <http://dx.doi.org/10.1111/j.1600-0838.2009.00949.x>
50. Kongsgaard M, Qvortrup K, Larsen J, et al. Fibril morphology and tendon mechanical properties in patellar tendinopathy: effects of heavy slow resistance training. *Am J Sports Med*. 2010;38:749-756. <http://dx.doi.org/10.1177/0363546509350915>
51. Kubo K, Kanehisa H, Fukunaga T. Effects of different duration isometric contractions on tendon elasticity in human quadriceps muscles. *J Physiol*. 2001;536:649-655. <http://dx.doi.org/10.1111/j.1469-7793.2001.0649c.xd>
52. Kujala UM, Sarna S, Kaprio J. Cumulative incidence of Achilles tendon rupture and tendinopathy in male former elite athletes. *Clin J Sport Med*. 2005;15:133-135. <http://dx.doi.org/10.1097/01.jsm.0000165347.55638.23>
53. Langberg H, Ellingsgaard H, Madsen T, et al. Eccentric rehabilitation exercise increases peritendinous type I collagen synthesis in humans with Achilles tendinosis. *Scand J Med Sci Sports*. 2007;17:61-66. <http://dx.doi.org/10.1111/j.1600-0838.2006.00522.x>
54. Langberg H, Olesen J, Skovgaard D, Kjaer M. Age related blood flow around the Achilles tendon during exercise in humans. *Eur J Appl Physiol*. 2001;84:246-248. <http://dx.doi.org/10.1007/s004210170013>
55. Langberg H, Skovgaard D, Petersen LJ, Bülow J, Kjaer M. Type I collagen synthesis and degradation in peritendinous tissue after exercise determined by microdialysis in humans. *J Physiol*. 1999;521 pt 1:299-306. <http://dx.doi.org/10.1111/j.1469-7793.1999.00299.x>
56. Lavagnino M, Arnoczky SP, Tian T, Vaupel Z. Effect of amplitude and frequency of cyclic tensile strain on the inhibition of MMP-1 mRNA expression in tendon cells: an in vitro study. *Connect Tissue Res*. 2003;44:181-187.
57. Lian ØB, Engebretsen L, Bahr R. Prevalence of jumper's knee among elite athletes from different sports: a cross-sectional study. *Am J Sports Med*. 2005;33:561-567. <http://dx.doi.org/10.1177/0363546504270454>
58. Lopes AD, Hespanhol Junior LC, Yeung SS, Costa LO. What are the main running-related musculoskeletal injuries? A systematic review. *Sports Med*. 2012;42:891-905. <http://dx.doi.org/10.1007/BF03262301>
59. Maeda E, Shelton JC, Bader DL, Lee DA. Time dependence of cyclic tensile strain on collagen production in tendon fascicles. *Biochem Biophys Res Commun*. 2007;362:399-404. <http://dx.doi.org/10.1016/j.bbrc.2007.08.029>
60. Maffulli N, Khan KM, Puddu G. Overuse tendon conditions: time to change a confusing terminology. *Arthroscopy*. 1998;14:840-843. [http://dx.doi.org/10.1016/S0749-8063\(98\)70021-0](http://dx.doi.org/10.1016/S0749-8063(98)70021-0)
61. Mafi N, Lorentzon R, Alfredson H. Superior short-term results with eccentric calf muscle training compared to concentric training in a randomized prospective multicenter study on patients with chronic Achilles tendinosis. *Knee Surg Sports Traumatol Arthrosc*. 2001;9:42-47. <http://dx.doi.org/10.1007/s001670000148>
62. Magnusson SP, Aagaard P, Dyhre-Poulsen P, Kjaer M. Load-displacement properties of the human triceps surae aponeurosis in vivo. *J Physiol*. 2001;531:277-288. <http://dx.doi.org/10.1111/j.1469-7793.2001.0277j.x>

63. Magnusson SP, Langberg H, Kjaer M. The pathogenesis of tendinopathy: balancing the response to loading. *Nat Rev Rheumatol*. 2010;6:262-268. <http://dx.doi.org/10.1038/nrrheum.2010.43>
64. Malliaras P, Barton CJ, Reeves ND, Langberg H. Achilles and patellar tendinopathy loading programmes: a systematic review comparing clinical outcomes and identifying potential mechanisms for effectiveness. *Sports Med*. 2013;43:267-286. <http://dx.doi.org/10.1007/s40279-013-0019-z>
65. Miller BF, Olesen JL, Hansen M, et al. Coordinated collagen and muscle protein synthesis in human patella tendon and quadriceps muscle after exercise. *J Physiol*. 2005;567:1021-1033. <http://dx.doi.org/10.1113/jphysiol.2005.093690>
66. Moerch L, Pingel J, Boesen M, Kjaer M, Langberg H. The effect of acute exercise on collagen turnover in human tendons: influence of prior immobilization period. *Eur J Appl Physiol*. 2013;113:449-455. <http://dx.doi.org/10.1007/s00421-012-2450-5>
67. Niesen-Vertommen SL, Taunton JE, Clement DB, Mosher RE. The effect of eccentric versus concentric exercise in the management of Achilles tendonitis. *Clin J Sport Med*. 1992;2:109-113. <http://dx.doi.org/10.1097/00042752-199204000-00006>
68. Nirmalanandhan VS, Shearn JT, Juncosa-Melvin N, et al. Improving linear stiffness of the cell-seeded collagen sponge constructs by varying the components of the mechanical stimulus. *Tissue Eng Part A*. 2008;14:1883-1891. <http://dx.doi.org/10.1089/ten.tea.2007.0125>
69. Nørregaard J, Larsen CC, Bieler T, Langberg H. Eccentric exercise in treatment of Achilles tendinopathy. *Scand J Med Sci Sports*. 2007;17:133-138. <http://dx.doi.org/10.1111/j.1600-0838.2006.00545.x>
70. Öhberg L, Alfredson H. Effects on neovascularisation behind the good results with eccentric training in chronic mid-portion Achilles tendinosis? *Knee Surg Sports Traumatol Arthrosc*. 2004;12:465-470. <http://dx.doi.org/10.1007/s00167-004-0494-8>
71. Öhberg L, Lorentzon R, Alfredson H. Eccentric training in patients with chronic Achilles tendinosis: normalised tendon structure and decreased thickness at follow up. *Br J Sports Med*. 2004;38:8-11; discussion 11. <http://dx.doi.org/10.1136/bjsm.2001.000284>
72. Parent G, Huppé N, Langelier E. Low stress tendon fatigue is a relatively rapid process in the context of overuse injuries. *Ann Biomed Eng*. 2011;39:1535-1545. <http://dx.doi.org/10.1007/s10439-011-0254-0>
73. Paulsen G, Mikkelsen UR, Raastad T, Peake JM. Leucocytes, cytokines and satellite cells: what role do they play in muscle damage and regeneration following eccentric exercise? *Exerc Immunol Rev*. 2012;18:42-97.
74. Pearson SJ, Burgess K, Onambele GN. Creep and the in vivo assessment of human patellar tendon mechanical properties. *Clin Biomech* (Bristol, Avon). 2007;22:712-717. <http://dx.doi.org/10.1016/j.clinbiomech.2007.02.006>
75. Pedersen BK, Steensberg A, Fischer C, et al. Searching for the exercise factor: is IL-6 a candidate? *J Muscle Res Cell Motil*. 2003;24:113-119.
76. Petersen W, Welp R, Rosenbaum D. Chronic Achilles tendinopathy: a prospective randomized study comparing the therapeutic effect of eccentric training, the AirHeel brace, and a combination of both. *Am J Sports Med*. 2007;35:1659-1667. <http://dx.doi.org/10.1177/0363546507303558>
77. Pingel J, Fredberg U, Mikkelsen LR, et al. No inflammatory gene-expression response to acute exercise in human Achilles tendinopathy. *Eur J Appl Physiol*. 2013;113:2101-2109. <http://dx.doi.org/10.1007/s00421-013-2638-3>
78. Purdam CR, Jonsson P, Alfredson H, Lorentzon R, Cook JL, Khan KM. A pilot study of the eccentric decline squat in the management of painful chronic patellar tendinopathy. *Br J Sports Med*. 2004;38:395-397. <http://dx.doi.org/10.1136/bjsm.2003.000053>
79. Rees JD, Lichtwark GA, Wolman RL, Wilson AM. The mechanism for efficacy of eccentric loading in Achilles tendon injury; an in vivo study in humans. *Rheumatology (Oxford)*. 2008;47:1493-1497. <http://dx.doi.org/10.1093/rheumatology/ken262>
80. Rees JD, Stride M, Scott A. Tendons – time to revisit inflammation. *Br J Sports Med*. 2014;48:1553-1557. <http://dx.doi.org/10.1136/bjsports-2012-091957>
81. Roche AJ, Calder JD. Achilles tendinopathy: a review of the current concepts of treatment. *Bone Joint J*. 2013;95-B:1299-1307. <http://dx.doi.org/10.1302/0301-620X.95B10.31881>
82. Rompe JD, Nafe B, Furia JP, Maffulli N. Eccentric loading, shock-wave treatment, or a wait-and-see policy for tendinopathy of the main body of tendo Achillis: a randomized controlled trial. *Am J Sports Med*. 2007;35:374-383. <http://dx.doi.org/10.1177/0363546506295940>
83. Roos EM, Engström M, Lagerquist A, Söderberg B. Clinical improvement after 6 weeks of eccentric exercise in patients with mid-portion Achilles tendinopathy – a randomized trial with 1-year follow-up. *Scand J Med Sci Sports*. 2004;14:286-295. <http://dx.doi.org/10.1111/j.1600-0838.2004.378.x>
84. Semmler JG, Tucker KJ, Allen TJ, Proske U. Eccentric exercise increases EMG amplitude and force fluctuations during submaximal contractions of elbow flexor muscles. *J Appl Physiol* (1985). 2007;103:979-989. <http://dx.doi.org/10.1152/jappphysiol.01310.2006>
85. Seynnes OR, Erskine RM, Maganaris CN, et al. Training-induced changes in structural and mechanical properties of the patellar tendon are related to muscle hypertrophy but not to strength gains. *J Appl Physiol* (1985). 2009;107:523-530. <http://dx.doi.org/10.1152/jappphysiol.00213.2009>
86. Shalabi A, Kristoffersen-Wilberg M, Svensson L, Aspelin P, Movin T. Eccentric training of the gastrocnemius-soleus complex in chronic Achilles tendinopathy results in decreased tendon volume and intratendinous signal as evaluated by MRI. *Am J Sports Med*. 2004;32:1286-1296. <http://dx.doi.org/10.1177/0363546504263148>
87. Shepherd JH, Screen HR. Fatigue loading of tendon. *Int J Exp Pathol*. 2013;94:260-270. <http://dx.doi.org/10.1111/iep.12037>
88. Silbernagel KG, Thomeé R, Eriksson BI, Karlsson J. Continued sports activity, using a pain-monitoring model, during rehabilitation in patients with Achilles tendinopathy: a randomized controlled study. *Am J Sports Med*. 2007;35:897-906. <http://dx.doi.org/10.1177/0363546506298279>
89. Silbernagel KG, Thomeé R, Thomeé P, Karlsson J. Eccentric overload training for patients with chronic Achilles tendon pain – a randomized controlled study with reliability testing of the evaluation methods. *Scand J Med Sci Sports*. 2001;11:197-206. <http://dx.doi.org/10.1034/j.1600-0838.2001.110402.x>
90. Stanish WD, Rubinovich RM, Curwin S. Eccentric exercise in chronic tendinitis. *Clin Orthop Relat Res*. 1986;65-68.
91. Stasinopoulos D, Stasinopoulos I. Comparison of effects of exercise programme, pulsed ultrasound and transverse friction in the treatment of chronic patellar tendinopathy. *Clin Rehabil*. 2004;18:347-352.
92. Sussmilch-Leitch SP, Collins NJ, Bialocerkowski AE, Warden SJ, Crossley KM. Physical therapies for Achilles tendinopathy: systematic review and meta-analysis. *J Foot Ankle Res*. 2012;5:15. <http://dx.doi.org/10.1186/1757-1146-5-15>
93. Tanaka S, Petersen M, Cameron L. Prevalence and risk factors of tendinitis and related disorders of the distal upper extremity among U.S. workers: comparison to carpal tunnel syndrome. *Am J Ind Med*. 2001;39:328-335. [http://dx.doi.org/10.1002/1097-0274\(200103\)39:3<328::AID-AJIM1021>3.0.CO;2-I](http://dx.doi.org/10.1002/1097-0274(200103)39:3<328::AID-AJIM1021>3.0.CO;2-I)
94. Toft AD, Jensen LB, Bruunsgaard H, et al. Cytokine response to eccentric exercise in young and elderly humans. *Am J Physiol Cell Physiol*. 2002;283:C289-C295. <http://dx.doi.org/10.1152/ajpcell.00583.2001>
95. Visnes H, Hoksrud A, Cook J, Bahr R. No effect of eccentric training on jumper's knee in volleyball players during the competitive season: a randomized clinical trial. *Clin J Sport Med*. 2005;15:227-234.
96. Wang JH. Mechanobiology of tendon. *J Biomech*. 2006;39:1563-1582. <http://dx.doi.org/10.1016/j.jbiomech.2005.05.011>
97. Webb K, Hitchcock RW, Smeal RM, Li W, Gray SD, Tresco PA. Cyclic strain increases fibroblast proliferation, matrix accumulation, and elastic modulus of fibroblast-seeded polyurethane constructs. *J Biomech*. 2006;39:1136-1144. <http://dx.doi.org/10.1016/j.jbiomech.2004.08.026>
98. Woo SL, Ritter MA, Amiel D, et al. The biomechanical and biochemical properties of swine tendons – long term effects of

exercise on the digital extensors. *Connect Tissue Res.* 1980;7:177-183. <http://dx.doi.org/10.3109/03008208009152109>

99. Yamamoto E, Hayashi K, Yamamoto N. Mechanical properties of collagen fascicles from stress-shielded patellar tendons in the rabbit. *Clin Biomech (Bristol, Avon)*. 1999;14:418-425.
100. Yamamoto E, Kogawa D, Tokura S, Hayashi K. Effects of the frequency and duration of cyclic stress on the mechanical properties of cultured collagen fascicles from the rabbit patellar tendon. *J Biomech Eng.* 2005;127:1168-1175.

101. Yang G, Crawford RC, Wang JH. Proliferation and collagen production of human patellar tendon fibroblasts in response to cyclic uniaxial stretching in serum-free conditions. *J Biomech.* 2004;37:1543-1550. <http://dx.doi.org/10.1016/j.jbiomech.2004.01.005>

102. Yelland MJ, Sweeting KR, Lyftogt JA, Ng SK, Scuffham PA, Evans KA. Prolotherapy injections and eccentric loading exercises for painful Achilles tendinosis: a randomised trial. *Br J Sports Med.* 2011;45:421-428. <http://dx.doi.org/10.1136/bjsm.2009.057968>

103. Young MA, Cook JL, Purdam CR, Kiss ZS, Alfredson H. Eccentric decline squat protocol offers superior results at 12 months compared with traditional eccentric protocol for patellar tendinopathy in volleyball players. *Br J Sports Med.* 2005;39:102-105. <http://dx.doi.org/10.1136/bjsm.2003.010587>



MORE INFORMATION
WWW.JOSPT.ORG

PUBLISH Your Manuscript in a Journal With International Reach

JOSPT offers authors of accepted papers an **international audience**. The *Journal* is currently distributed to the members of APTA's Orthopaedic and Sports Physical Therapy Sections and **33 orthopaedics, manual therapy, and sports groups in 26 countries** who provide online access either as a member benefit or at a discount. As a result, the *Journal* is now distributed monthly to more than **30,000 individuals around the world** who specialize in musculoskeletal and sports-related rehabilitation, health, and wellness. In addition, *JOSPT* reaches students and faculty, physical therapists and physicians at more than **1,500 institutions in 56 countries**.

Please review our Information for and Instructions to Authors at www.jospt.org in the **Info Center** for **Authors** and submit your manuscript for peer review at <http://mc.manuscriptcentral.com/jospt>.